By: Namazzi Christine Beepath Clinical and Counselling Psychologist azzi@gmail.com Tel. +256 773185696 cnamazzi@gmail.com

SCOPE OF COVERAGE


TRAUMA = EVENT+ EXPERIENCE OF THE EVENT+EFFECTS

Trauma can be Acute. Chronic, and complex or attachment trauma

1-INTRODUCTION

- A global survey found that among 68.894 people from 24 countries, 70.4% experienced lifetime traumas. Around 76 million women worldwide suffer from substance use disorders (SUDs), with
- In East Africa, women's alcohol consumption rates are increasing, with Kenya reporting 21.2% of women consuming alcohol, and **Uganda** showing
- In Uganda, cannabis comes next to alcohol among common illicit drugs used by women, with 2.5% of the female population using it annually.

4-THE INTERTWINE BETWEEN TRAUMA AND ALCOHOL USE DISORDER "GREAT IMPACT"



2-RESULTS AND FINDINGS

trauma specific models in Uganda is almost

3-AIMS/OBJECTIVES AND PURPOSE

The present presentation is to assess the

INTERPERSONAL TRAUMA





5-TREATMENT


The principles of Trauma Informed Care

The 4 R's of Trauma informed care:

- Recognise,Respond andResist re-traumatisation.

policy, therapeutic parenting skills, trauma sensitive approaches, attachment-based therapy, improvement of maternal functionality and teach self regulation

6-LOCATION





Kawotto-Kajjansi Wafula Road

Albert Elwa Louis



7-CONTACTS

Mental health care providers should be sensitised more on the relationship between AUD and Trauma especially among women who come for treatment. How the intertwine between the two can affect treatment outcomes.

Trauma informed and trauma responsive care is found to be effective in AUD treatment among women,

Ministry of Health and different organisations looking at prevention and treatment of AUD such as UAPA and APRAU should encourage treatment centres to incorporate trauma informed care and trauma specific models into the services provided for effective AUD treatment and sustainability of recovery among women.

